

Lesson 4 Practice Problems

1. If the equation $(x + 10)x = 0$ is true, which statement is also true according to the zero product property?

- A. only $x = 0$
- B. either $x = 0$ or $x + 10 = 0$
- C. either $x^2 = 0$ or $10x = 0$
- D. only $x + 10 = 0$

2. What are the solutions to the equation $(10 - x)(3x - 9) = 0$?

- A. -10 and 3
- B. -10 and 9
- C. 10 and 3
- D. 10 and 9

3. Solve each equation.

- a. $(x - 6)(x + 5) = 0$
- b. $(x - 3)\left(\frac{2}{3}x - 6\right) = 0$
- c. $(-3x - 15)(x + 7) = 0$

4. Consider the expressions $(x - 4)(3x - 6)$ and $3x^2 - 18x + 24$.

Show that the two expressions define the same function.

5. Kiran saw that if the equation $(x + 2)(x - 4) = 0$ is true, then, by the zero product property, either $x + 2$ is 0 or $x - 4$ is 0. He then reasoned that, if $(x + 2)(x - 4) = 72$ is true, then either $x + 2$ is equal to 72 or $x - 4$ is equal to 72.

Explain why Kiran's conclusion is incorrect.

6. Andre wants to solve the equation $5x^2 - 4x - 18 = 20$. He uses a graphing calculator to graph $y = 5x^2 - 4x - 18$ and $y = 20$ and finds that the graphs cross at the points $(-2.39, 20)$ and $(3.19, 20)$.

a. Substitute each x -value Andre found into the expression $5x^2 - 4x - 18$. Then evaluate the expression.

b. Why did neither solution make $5x^2 - 4x - 18$ equal exactly 20?

(From Unit 7, Lesson 2.)

7. Select **all** the solutions to the equation $7x^2 = 343$.

A. 49

B. $-\sqrt{7}$

C. 7

D. -7

E. $\sqrt{49}$


F. $\sqrt{-49}$

G. $-\sqrt{49}$


(From Unit 7, Lesson 3.)

8. Here are two graphs that correspond to two patients, A and B. Each graph shows the amount of insulin, in micrograms (mcg) in a patient's body h hours after receiving an injection. The amount of insulin in each patient decreases exponentially.

Patient A



Patient B




Select **all** statements that are true about the insulin level of the two patients.

- A. After the injection, the patients have the same amount of insulin in their bodies.
- B. An equation for the micrograms of insulin, a , in Patient A's body h hours after the injection is $a = 200 \cdot \left(\frac{3}{5}\right)^h$.
- C. The insulin in Patient A is decaying at a faster rate than in Patient B.
- D. After 3 hours, Patient A has more insulin in their body than Patient B.
- E. At some time between 2 and 3 hours, the patients have the same insulin level.

(From Unit 5, Lesson 6.)

9. Han says this pattern of dots can be represented by a quadratic relationship because the dots are arranged in a rectangle in each step.

Do you agree? Explain your reasoning.



(From Unit 6, Lesson 2.)