

## Lesson 14: Solving Exponential Equations

- Let's solve equations using logarithms.

### 14.1: A Valid Solution?

To solve the equation  $5 \cdot e^{3a} = 90$ , Lin wrote the following:

$$\begin{aligned}5 \cdot e^{3a} &= 90 \\e^{3a} &= 18 \\3a &= \log_e 18 \\a &= \frac{\log_e 18}{3}\end{aligned}$$

Is her solution valid? Be prepared to explain what she did in each step to support your answer.

## 14.2: Natural Logarithm

1. Complete the table with equivalent equations. The first row is completed for you.

	exponential form	logarithmic form
a.	$e^0 = 1$	$\ln 1 = 0$
b.	$e^1 = e$	
c.	$e^{-1} = \frac{1}{e}$	
d.		$\ln \frac{1}{e^2} = -2$
e.	$e^x = 10$	

2. Solve each equation by expressing the solution using  $\ln$  notation. Then, find the approximate value of the solution using the “ln” button on a calculator.

a.  $e^m = 20$

b.  $e^n = 30$

c.  $e^p = 7.5$

## 14.3: Solving Exponential Equations

Without using a calculator, solve each equation. It is expected that some solutions will be expressed using log notation. Be prepared to explain your reasoning.

1.  $10^x = 10,000$

2.  $5 \cdot 10^x = 500$

3.  $10^{(x+3)} = 10,000$

4.  $10^{2x} = 10,000$

5.  $10^x = 315$

6.  $2 \cdot 10^x = 800$

7.  $10^{(1.2x)} = 4,000$

8.  $7 \cdot 10^{(0.5x)} = 70$

9.  $2 \cdot e^x = 16$

10.  $10 \cdot e^{3x} = 250$

### Are you ready for more?

1. Solve the equations  $10^n = 16$  and  $10^n = 2$ . Express your answers as logarithms.
2. What is the relationship between these two solutions? Explain how you know.

### Lesson 14 Summary

So far we have solved exponential equations by

- finding whole number powers of the base (for example, the solution of  $10^x = 100,000$  is 5)
- estimation (for example, the solution of  $10^x = 300$  is between 2 and 3)
- using a logarithm and approximating its value on a calculator (for example, the solution of  $10^x = 300$  is  $\log 300 \approx 2.48$ )

Sometimes solving exponential equations takes additional reasoning. Here are a couple of examples.

$$\begin{aligned} 5 \cdot 10^x &= 45 \\ 5 \cdot 10^x &= 45 \\ 10^x &= 9 \\ x &= \log 9 \end{aligned}$$

$$\begin{aligned} 10^{(0.2t)} &= 1,000 \\ 10^{(0.2t)} &= 10^3 \\ 0.2t &= 3 \\ t &= \frac{3}{0.2} \\ t &= 15 \end{aligned}$$

In the first example, the power of 10 is multiplied by 5, so to find the value of  $x$  that makes this equation true each side was divided by 5. From there, the equation was rewritten as a logarithm, giving an exact value for  $x$ .

In the second example, the expressions on each side of the equation were rewritten as powers of 10:  $10^{(0.2t)} = 10^3$ . This means that the exponent  $0.2t$  on one side and the 3 on

the other side must be equal, and leads to a simpler expression to solve where we don't need to use a logarithm.

How do we solve an exponential equation with base  $e$ , such as  $e^x = 5$ ? We can express the solution using the **natural logarithm**, the logarithm for base  $e$ . Natural logarithm is written as  $\ln$ , or sometimes as  $\log_e$ . Just like the equation  $10^2 = 100$  can be rewritten, in logarithmic form, as  $\log_{10} 100 = 2$ , the equation  $e^0 = 1$  can be rewritten as  $\ln 1 = 0$ . Similarly,  $e^{-2} = \frac{1}{e^2}$  can be rewritten as  $\ln \frac{1}{e^2} = -2$ .

All this means that we can solve  $e^x = 5$  by rewriting the equation as  $x = \ln 5$ . This says that  $x$  is the exponent to which base  $e$  is raised to equal 5.

To estimate the size of  $\ln 5$ , remember that  $e$  is about 2.7. Because 5 is greater than  $e^1$ , this means that  $\ln 5$  is greater than 1.  $e^2$  is about  $(2.7)^2$  or 7.3. Because 5 is less than  $e^2$ , this means that  $\ln 5$  is less than 2. This suggests that  $\ln 5$  is between 1 and 2. Using a calculator we can check that  $\ln 5 \approx 1.61$ .