# Lesson 5: How Many Groups? (Part 2)

## 5.1: Reasoning with Fraction Strips

Write a fraction or whole number as an answer for each question. If you get stuck, use the fraction strips. Be prepared to share your reasoning.

1. How many $\frac{1}{2}$s are in 2?

2. How many $\frac{1}{5}$s are in 3?

3. How many $\frac{1}{8}$s are in $1\frac{1}{4}$?

4. $1 \div \frac{2}{6} = ?$

5. $2 \div \frac{2}{9} = ?$

6. $4 \div \frac{2}{10} = ?$

| 1 | | | 1 | | |
|---|---|---|---|---|---|
| $\frac{1}{2}$ | | $\frac{1}{2}$ | $\frac{1}{2}$ | | $\frac{1}{2}$ |
| $\frac{1}{3}$ | $\frac{1}{3}$ | $\frac{1}{3}$ | $\frac{1}{3}$ | $\frac{1}{3}$ | $\frac{1}{3}$ |
| $\frac{1}{4}$ $\frac{1}{4}$ | $\frac{1}{4}$ | $\frac{1}{4}$ | $\frac{1}{4}$ | $\frac{1}{4}$ | $\frac{1}{4}$ $\frac{1}{4}$ |
| $\frac{1}{5}$ $\frac{1}{5}$ | $\frac{1}{5}$ | $\frac{1}{5}$ $\frac{1}{5}$ | $\frac{1}{5}$ $\frac{1}{5}$ | $\frac{1}{5}$ | $\frac{1}{5}$ $\frac{1}{5}$ |
| $\frac{1}{6}$ $\frac{1}{6}$ | $\frac{1}{6}$ $\frac{1}{6}$ | $\frac{1}{6}$ $\frac{1}{6}$ | $\frac{1}{6}$ $\frac{1}{6}$ | $\frac{1}{6}$ $\frac{1}{6}$ | $\frac{1}{6}$ $\frac{1}{6}$ |
| $\frac{1}{8}$ $\frac{1}{8}$ | $\frac{1}{8}$ $\frac{1}{8}$ | $\frac{1}{8}$ $\frac{1}{8}$ | $\frac{1}{8}$ $\frac{1}{8}$ | $\frac{1}{8}$ $\frac{1}{8}$ | $\frac{1}{8}$ $\frac{1}{8}$ |
| $\frac{1}{9}$ $\frac{1}{9}$ $\frac{1}{9}$ | $\frac{1}{9}$ $\frac{1}{9}$ | $\frac{1}{9}$ $\frac{1}{9}$ | $\frac{1}{9}$ $\frac{1}{9}$ | $\frac{1}{9}$ $\frac{1}{9}$ | $\frac{1}{9}$ $\frac{1}{9}$ $\frac{1}{9}$ |

## 5.2: More Reasoning with Pattern Blocks

Your teacher will give you pattern blocks. Use them to answer the questions.

1. If the trapezoid represents 1 whole, what do each of the other shapes represent? Be prepared to show or explain your reasoning.



2. Use pattern blocks to represent each multiplication equation. Use the trapezoid to represent 1 whole.

   a. $3 \cdot \frac{1}{3} = 1$

   b. $3 \cdot \frac{2}{3} = 2$

3. Diego and Jada were asked "How many rhombuses are in a trapezoid?"

   ○ Diego says, "$1\frac{1}{3}$. If I put 1 rhombus on a trapezoid, the leftover shape is a triangle, which is $\frac{1}{3}$ of the trapezoid."

   ○ Jada says, "I think it's $1\frac{1}{2}$. Since we want to find out 'how many rhombuses,' we should compare the leftover triangle to a rhombus. A triangle is $\frac{1}{2}$ of a rhombus."

   Do you agree with either of them? Explain or show your reasoning.

**Grade 6 Unit 4**
Lesson 5

CC BY Open Up Resources. Adaptations CC BY IM.

2

4. Select **all** the equations that can be used to answer the question: "How many rhombuses are in a trapezoid?"

- $\frac{2}{3} \div ? = 1$
- $1 \div \frac{2}{3} = ?$
- $? \div \frac{2}{3} = 1$
- $? \cdot \frac{2}{3} = 1$
- $1 \cdot \frac{2}{3} = ?$

## 5.3: Drawing Diagrams to Show Equal-sized Groups

For each situation, draw a diagram for the relationship of the quantities to help you answer the question. Then write a multiplication equation or a division equation for the relationship. Be prepared to share your reasoning.

1. The distance around a park is $\frac{3}{2}$ miles. Noah rode his bicycle around the park for a total of 3 miles. How many times around the park did he ride?

2. You need $\frac{3}{4}$ yard of ribbon for one gift box. You have 3 yards of ribbon. How many gift boxes do you have ribbon for?

3. The water hose fills a bucket at $\frac{1}{3}$ gallon per minute. How many minutes does it take to fill a 2-gallon bucket?

### Are you ready for more?

How many heaping teaspoons are in a heaping tablespoon? How would the answer depend on the shape of the spoons?

**Grade 6 Unit 4**
Lesson 5

CC BY Open Up Resources. Adaptations CC BY IM.

3

## Lesson 5 Summary

Suppose one batch of cookies requires $\frac{2}{3}$ cup flour. How many batches can be made with 4 cups of flour?

We can think of the question as being: "How many $\frac{2}{3}$ are in 4?" and represent it using multiplication and division equations.

$$? \cdot \frac{2}{3} = 4$$

$$4 \div \frac{2}{3} = ?$$

Let's use pattern blocks to visualize the situation and say that a hexagon is 1 whole.



Since 3 rhombuses make a hexagon, 1 rhombus represents $\frac{1}{3}$ and 2 rhombuses represent $\frac{2}{3}$. We can see that 6 pairs of rhombuses make 4 hexagons, so there are 6 groups of $\frac{2}{3}$ in 4.

Other kinds of diagrams can also help us reason about equal-sized groups involving fractions. This example shows how we might reason about the same question from above: "How many $\frac{2}{3}$-cups are in 4 cups?"



We can see each "cup" partitioned into thirds, and that there are 6 groups of $\frac{2}{3}$-cup in 4 cups. In both diagrams, we see that the unknown value (or the "?" in the equations) is 6. So we can now write:

$$6 \cdot \frac{2}{3} = 4$$

$$4 \div \frac{2}{3} = 6$$

**Grade 6 Unit 4**

Lesson 5

CC BY Open Up Resources. Adaptations CC BY IM.

4